

Physical Effects of Chemically & Physically Dispersed Oil on Wildlife

This study examines the effects of a marine bird or otter diving through a sub-surface plume of dispersed oil.


The Physical Effects of Chemically and Physically Dispersed Oil on Wildlife



Rebecca S. Duerr
Michael H. Ziccardi
J. Gregory Massey
Yvonne N. Addassi

Chemical Dispersants

- Surfactants or solvents that disperse oil into water column
- Applied by aircraft spraying
- Target dilution 1:10 to 1:60 dispersant:oil
- Maximum oil concentration in solution in water column = ~400 ppm



Oil Spill Dispersants: Efficacy and Effects 2005

Tradeoffs of dispersant use

Pro

- Reduce potential impact on water surface wildlife
- Prevent slick reaching sensitive coastal areas

Con


- Increased direct toxicity to water column and benthic organisms
- Limited time frame of utility
- Long term storage and maintenance of ready state over years between events

Study rationale


- Pre-approval partly based on assumed benefit to surface vertebrate wildlife
- Limited previous studies of dispersed oil on vertebrate wildlife
- Direct exposure of dispersants to animals is inevitable with application



But first: a Feather anatomy primer

Generic contour feather











Redrawn after Lucas and Stettenheim 1972.

Proctor and Lynch 1993





Hypotheses

1. Common murre contour feathers exposed to sea water mixed with mechanically dispersed oil, chemically dispersed oil, or dispersant (only) will display significant and dose-responsive alterations in microscopic cohesion of barbs, barbules and hooklets as compared among treatment groups and as compared to control feathers exposed to plain sea water.

Hypotheses


2. Sea otter hair similarly exposed will display visual alteration in structure as compared to control hair exposed to plain sea water
3. Quantified total petroleum hydrocarbon (TPH) levels will differ significantly among treatment groups and will display a dose-response relationship to increasing exposure concentrations of oil.


Methods


- Test materials:
 - Alaska North Slope Crude Oil
 - Corexit 9500 (Nalco-Exxon) dispersant
 - Instant Ocean (artificial sea water)
- Test solutions mixed per CROSERF specs

Treatment groups


	Control	Dispersant t only	Oil only	Dispersant + oil
[Low] = 40 ppm	0 Disp + 0 oil/L	2 μ L Disp + 0 oil/L	0 Disp + 44 μ L oil/L	2 μ L Disp + 44 μ L oil/L
[Mod] = 200 ppm		11 μ L Disp + 0 oil/L	0 Disp + 222 μ L oil/L	11 μ L Disp + 222 μ L oil/L
[High] = 400 ppm		22 μ L Disp + 0 oil/L	0 Disp + 444 μ L oil/L	22 μ L Disp + 444 μ L oil/L









100X





100X




400X




1000X

1 cm central
area for
microscopy




4.0 cm total length







Clean feather 100X




Clean feather 400X




Clean feather 1000X



100% oil
100X





100% oil
400X

$\sin A = 50/61.5$
 $= 54.4^\circ$

A

131 μm

61.5 μm

50 μm

14 μm

131 μm

17.1 μm

11.3 μm

14.7 μm

13.8 μm

11.8 μm

13.8 μm

100% oil
400X

Project Timeline

- June: exposure trials and sample photography
- July-Aug: photographic and numeric data analysis

Future Directions

- Effects of exposure on penetration pressure of whole pieces of pelt
- Effects of surface tension on barbule friction and air bubble formation
- 3D relationship of adjacent feathers

References

- Addassi, Y., M. Sowby, H. Parker-Hall and W. Robberson. 2005a. Establishment of Dispersant Use Zones in the State of California: A consensus approach form marine waters 3 – 200 nautical miles from shore. *Proceedings of the 2005 International Oil Spill Conference*, Miami Beach, Florida. American Petroleum Institute, Washington, D.C.
- Addassi., Y., E. Faurot-Daniels. 2005b. California Oil Spill Dispersant Plan – Achievement Through Cooperation . *Proceedings of the 2005 International Oil Spill Conference*, Miami Beach, Florida. American Petroleum Institute, Washington, D.C.
- Aurand, D., G.M. Coelho, R.G. Pond, J. Kraly, B. Martin, M. Sowby, J. Caplis and A.H. Walker. 2001. Results from cooperative ecological risk assessments for oil spill response planning in Galveston Bay, Texas and the San Francisco Bay area, California. 167- 175 In *Proceedings of the 2001 International Oil Spill Conference*, March 25-29, Tampa, Florida. American Petroleum Institute, Washington DC.
- Kraly, J., R.G. Pond, D.V. Aurand, G.M. Coelho, A.H. Walker, B. Martin, J. Caplis and M. Sowby. 2001. Ecological risk assessment principles applied to oil spill response planning. Pp 177-183 in *Proceeding of the 2001 International Oil Spill Conference*, Tampa, Florida. American Petroleum Institute, Washington, D.C.
- National Research Council (NRC) 1989, *Using Oil Spill Dispersants on the Sea*. National Academy Press, Washington, D.C.
- National Research Council (NRC) 2005. *Oil Spill Dispersants: Efficacy and Effects*. National Academy Press, Washington D.C.
- Pond, R.G. Aurand, and J.A. Kraly (compilers). 2000. Ecological Risk Assessment principles applied to oil spill response planning in the San Francisco Bay area. California Office of Spill Prevention and Response, Sacramento, CA. 70 + App.

Questions?

